



Hormones

VETEGLAN

Prevention and treatment of MMA
Parturition regulator

PHARMACEUTICAL FORM:

Injectable solution

COMPOSITION:

d-Cloprostenol 0,075 mg/ml



INDICATIONS:

Cows: Silent heats. Induction and synchronisation of oestrus. Luteinised cysts and persistent corpus luteum. Coadjutant treatment in pyometra and endometritis. Expulsion of mummified foetus. Induction of abortion at the first stage of gestation. Induction of calving.

Sows: Induction of farrowing. **Mares:** Treatment of persistent corpus luteum. Induction of oestrus.

DOSAGE AND ADMINISTRATION ROUTE:

Veteglan® should be administered by intramuscular route.

Cows: 150 mg/animal (equivalent to 2 ml VETEGLAN/animal). **Induction of oestrus:** administer VETEGLAN after establishing the existence of a functional corpus luteum (dosage of progesterone) generally between day 6 and 18 of the oestral cycle. The ovulation appears between 48 to 60 hours after treatment. If there is no sign of oestrus, the treatment may be repeated 11 days after the first treatment. **Induction of calving:** Administer VETEGLAN after the 270th day of gestation. Most animals are expected to respond within a period of 30-60 hours following injection. **Synchronisation of oestrus:** administration of two injections at an interval of 11 days. **Pyometra, endometria or mummified foetus expulsion:** One injection of VETEGLAN usually results in the expulsion of the mummified foetus from the uterus. When necessary, treatment may be repeated after 10 days. **Abortion:** Unwanted pregnancies can be efficiently terminated with the injection of VETEGLAN on the first 4 months of gestation. **Presence of persistent corpus luteum:** Administer VETEGLAN and observe the appearance of the next oestrus.

Sows: 75 µg/animal (equivalent to 1 ml VETEGLAN /animal). **Induction of farrowing:** Administer VETEGLAN from the 112th day of gestation, 70% of the animals is expected to respond within a period of 20 hours following injection. **Mares:** 75 µg/animal (equivalent to 1 ml VETEGLAN /animal). **Induction of oestrus:** Administer VETEGLAN after verifying the presence of a functional corpus luteum (dosage of progesterone). **Induction of foaling:** Administer VETEGLAN from the 320th day of gestation, most animals are expected to respond within few hours following injection. **Early foetal death:** Administer VETEGLAN after detecting the presence of an active corpus luteum (dosage of progesterone).

CONTRAINDICATIONS:

Do not administer to pregnant animals unless induced abortion is desired.

WITHDRAWAL PERIOD: Not applicable

REG. N°: 1798-ESP

PRESENTATION : 20 ml.